



# Correction for Khan et al., “A Multicenter Study To Evaluate Ceftaroline Breakpoints: Performance in an Area with High Prevalence of Methicillin-Resistant *Staphylococcus aureus* Sequence Type 5 Lineage”

Ayesha Khan,<sup>a,b,c</sup> Lina M. Rivas,<sup>d,e</sup> Maria Spencer,<sup>d,e</sup> Rodrigo Martinez,<sup>d,e</sup> Marusella Lam,<sup>f</sup> Pamela Rojas,<sup>g</sup> Lorena Porte,<sup>d,e</sup> Francisco Silva,<sup>h</sup> Stephanie Braun,<sup>i</sup> Francisca Valdivieso,<sup>j</sup> Margareta Mülhauser,<sup>k</sup> Mónica Lafourcade,<sup>l</sup> William R. Miller,<sup>a</sup> Patricia García,<sup>e,f</sup> Cesar A. Arias,<sup>a,b,e,m,n</sup> Jose M. Munita<sup>a,d,e</sup>

<sup>a</sup>Center for Antimicrobial Resistance and Microbial Genomics and Division of Infectious Diseases, University of Texas Health Science Center, McGovern Medical School, Houston, Texas, USA

<sup>b</sup>Department of Microbiology and Molecular Genetics, University of Texas Health Science Center, McGovern Medical School, Houston, Texas, USA

<sup>c</sup>MD Anderson Cancer Center, UT Health Graduate School of Biomedical Sciences, Houston, Texas, USA

<sup>d</sup>Genomics and Resistant Microbes Group, Facultad de Medicina-Clinica Alemana, Universidad del Desarrollo, Santiago, Chile

<sup>e</sup>Millennium Initiative for Collaborative Research on Bacterial Resistance (MICROB-R), Millennium Science Initiative, Santiago, Chile

<sup>f</sup>Departamento de Laboratorios Clínicos, Escuela de Medicina, Pontificia Universidad Católica de Chile, Santiago, Chile

<sup>g</sup>Hospital Padre Hurtado, Santiago, Chile

<sup>h</sup>Hospital Clínico Universidad de Chile, Santiago, Chile

<sup>i</sup>Hospital Militar, Santiago, Chile

<sup>j</sup>Hospital Dr. Luis Calvo Mackenna, Santiago, Chile

<sup>k</sup>Hospital Dipreca, Santiago, Chile

<sup>l</sup>Clinica Santa María, Santiago, Chile

<sup>m</sup>Molecular Genetics and Antimicrobial Resistance Unit, International Center for Microbial Genomics, Universidad El Bosque, Bogotá, Colombia

<sup>n</sup>Center for Infectious Diseases, University of Texas Health Science Center, School of Public Health, Houston, Texas, USA

Volume 57, issue 9, e00798-19, 2019, <https://doi.org/10.1128/JCM.00798-19>. The following funding information and acknowledgments should have appeared in the article.

Jose M. Munita was supported by Comision Nacional de Investigacion Cientifica y Tecnologica (CONICYT) (grant number FONDECYT 1171805) and the ANID Millenium Science Initiative, MICROB-R, NCN17\_081, Government of Chile. Cesar A. Arias was supported by the National Institute of Allergy and Infectious Diseases (NIAID) (grant numbers K24-AI121296, R01AI093749, R01AI134637, and R21/R33 AI121519).

**Citation** Khan A, Rivas LM, Spencer M, Martinez R, Lam M, Rojas P, Porte L, Silva F, Braun S, Valdivieso F, Mülhauser M, Lafourcade M, Miller WR, García P, Arias CA, Munita JM. 2021.

Correction for Khan et al., “A multicenter study to evaluate ceftaroline breakpoints: performance in an area with high prevalence of methicillin-resistant *Staphylococcus aureus* sequence type 5 lineage.” J Clin Microbiol 59:e00734-21. <https://doi.org/10.1128/JCM.00734-21>.

**Copyright** © 2021 American Society for Microbiology. All Rights Reserved.

Address correspondence to Cesar A. Arias, Cesari.Arias@uth.tmc.edu, or Jose M. Munita, Jose.M.Munita@uth.tmc.edu.

**Published** 19 May 2021