

Biphasic effect of linoleic acid on connexin 46 hemichannels

Mauricio A. Retamal · Flavio Evangelista-Martínez ·
Carmen G. León-Paravic · Guillermo A. Altenberg ·
Luis Reuss

Received: 10 November 2010 / Revised: 1 February 2011 / Accepted: 4 February 2011 / Published online: 1 March 2011
© Springer-Verlag 2011

Abstract Connexins form hemichannels at undocked plasma membranes and gap-junction channels (GJs) at intercellular contacting zones. Under physiological conditions, hemichannels have low open probabilities, but their activation under pathological conditions, such as ischemia, induces and/or accelerates cell death. Connexin 46 (Cx46) is a major connexin of the lens, and mutations of this connexin induce cataracts. Here, we report the effects of linoleic acid (LA) on the electrical properties of Cx46 GJs and hemichannels expressed in *Xenopus laevis* oocytes. LA has a biphasic effect, increasing hemichannel current at 0.1 μ M and decreasing it at concentrations of 100 μ M or higher. The effects of extracellular and microinjected LA conjugated to coenzyme A (LA-CoA) suggest that the current activation site is accessible from the intracellular but not extracellular compartment, whereas the current inhibitory site is either located in a region of the hemichannel pore inaccessible to intracellular LA-CoA, or requires crossing of LA through an organelle membrane. Experiments with other fatty acids demonstrated that the block of hemichannels depends on the presence of a hydrogenated double bond at position 9 and is directly proportional to the number of double bonds. Experiments in paired oocytes expressing Cx46 showed that

LA does not affect GJs. The block by unsaturated fatty acids reported here opens the possibility that increases in the concentration of these lipids in the lens induce cataract formation by blocking Cx46 hemichannels.

Keywords Gap junction · Cataract · Frog oocyte · Lens fiber · Cx46 · Ion channel

Introduction

A gap-junction channel (GJC) is formed by head-to-head docking of two hemichannels, one from each adjacent cell. These hemichannels, also named connexons, are hexamers of proteins called connexins. The presence of undocked hemichannels at the plasma membrane has been demonstrated in several cell types, including lens fibers [16, 50]. However, their role under physiological or pathological conditions is not well defined, although several reports support the idea that controlled hemichannel opening allows autocrine/paracrine cell communication. For example, it has been demonstrated that hemichannels are involved in the release of signaling molecules such as ATP [54], glutamate [58], NAD^+ [10], and prostaglandin E2 [12], as well as in the uptake of glucose [46]. In pathological conditions such as inhibition of metabolism, hemichannels contribute to cell damage because their opening induces or accelerates cell death due to loss of metabolites and ion gradients, and Ca^{2+} entry [5, 50, 51, 55].

Lacking a vascular system, lens fibers use GJC as a nutritional and excretory pathway, where the flows of ions, water, and metabolites between the cells are determined by their chemical or electrochemical gradients [23, 38]. At least three connexin isoforms are expressed in the lens: Cx43 [6], Cx46 [45], and Cx50 [32], and it has been recently proposed that Cx46 and Cx50 hemichannels play a role in Na^+ and

M. A. Retamal (✉) · F. Evangelista-Martínez ·
C. G. León-Paravic
Laboratorio de Fisiología, Facultad de Medicina, Clínica
Alemana-Universidad del Desarrollo,
Avenida Las Condes 12438,
Santiago, Chile
e-mail: mretamal@udd.cl

M. A. Retamal · G. A. Altenberg · L. Reuss
Department of Cell Physiology and Molecular Biophysics
and Center for Membrane Protein Research,
Texas Tech University Health Sciences Center,
Lubbock, TX 79430-6551, USA

Ca^{2+} influx in lens fibers [16]. The importance of connexins in cataract formation is well documented, with Cx46- and Cx50-knockout mice developing cataracts at early ages, suggesting that connexin function is essential to maintain lens transparency [22, 43]. Increased levels of unsaturated fatty acids in the lens have been shown to induce and/or accelerate the development of cataracts [27, 33], but the mechanism of the damage of lens fibers is unknown. Although generation of radicals and lipid peroxidation products likely plays a role, they are not sufficient to explain the cytotoxicity [see 27]. Linoleic acid (LA), an unsaturated omega-6 fatty acid essential for biosynthesis of arachidonic acid [53], induces cataract formation in human [37] and bovine lens [21, 40]. Although there is no information on lens connexin isoforms, some isoforms are known to be sensitive to fatty acids. For example, oleamide-derived molecules inhibit Cx26 GJC [41].

Since Cx46 is essential for normal lens function, fatty acids are known to modulate connexins, and LA induces cataract formation, we decided to test whether LA alters the function of Cx46, one of the major connexins in the lens. In two-electrode voltage-clamp studies, we found a biphasic effect of LA on Cx46 hemichannels: low concentrations increased Cx46 hemichannel currents, whereas higher concentrations decreased them. We also observed a direct correlation between the number of fatty acid double bonds and the magnitude of the hemichannel current block, and found that a double bond at C9 is essential for hemichannel current inhibition. Unexpectedly, LA at low or high concentration did not affect Cx46 GJC.

Methods

Chemicals

Linoleic acid, palmitic acid, linoleoyl-CoA, and calphostin C were purchased from Sigma-Aldrich (Schnelldorf, Germany). Oleic acid, arachidonic acid, 9-thiastearic acid, and 9-nitrooleate were purchased from Cayman (Ann Arbor, Michigan, USA), and BAPTA-AM from Calbiochem (San Diego, California, USA).

Plasmid engineering, cRNA preparation, and injection into *Xenopus laevis* oocytes

The plasmid pSP64T-Cx46 containing rat Cx46 DNA was obtained from Dr. Lisa Ebihara (Rosalind Franklin University) [45]. Oocytes were injected with 12.5 ng of antisense Cx38 oligonucleotide alone or in combination with 25 ng of cRNA coding for Cx46. After cRNA injection, oocytes were maintained in Barth's solution (88 mM NaCl, 1 mM KCl, 5 mM CaCl_2 , 0.8 mM MgCl_2 , and 10 mM HEPES/

NaOH; pH 7.4, supplemented with 0.1 mg/ml gentamycin and 20 U/ml penicillin–streptomycin each) for 24–48 h, to allow for a good expression level. Additional details have been published [2].

Electrophysiological recordings and calculations

Whole-cell hemichannel currents were measured as described [47]. Briefly, oocytes were placed in a 1-ml recording chamber and superfused with ND96 solution (96 mM NaCl, 2 mM KCl, 1.8 mM CaCl_2 , and 5 mM HEPES/NaOH; pH 7.4) at room temperature. Glass microelectrodes were filled with 3 M KCl and had tip resistances of 0.5–1.5 M Ω when immersed in ND96. For data acquisition and analysis, we used a voltage-clamp amplifier (OC-725C, Warner Instruments, Hamden, CT) with pCLAMP 10 and a Digidata 1440A A/D Board (Molecular Devices, Foster City, CA). Currents were measured following 15-s rectangular voltage pulses, ranging from –50 to +60 mV, in 10-mV steps, with a holding potential of –60 mV and 10-s intervals between pulses. Current–voltage (I – V) relationships were calculated from the peak current values. Currents through GJCs were measured in paired oocytes. Briefly, both cells of the pair were clamped at –40 mV, and junctional currents were measured after changing the cell-membrane voltage of one cell to values between –140 and +60 mV (20-mV steps, 15-s intervals between pulses), while holding constant the voltage of the other cell (used as reference). The current supplied to the cell clamped at –40 mV is equal in amplitude, but opposite in sign, to the transjunctional current. To evaluate the effects of fatty acids, 500 μl of lipids dissolved in ND96 was carefully added to the 1-ml recording chamber, to final concentrations ranging from 0.1 to 1,000 μM . Unless otherwise indicated, the recordings were performed after 5 min of incubation with the fatty acids

Statistical analysis

Results are expressed as means \pm SEM, and n refers to the number of independent experiments. For statistical analysis, each treatment was compared to its respective control, and significance was determined using a one-way ANOVA or paired Student's t test, as appropriate. Differences were considered significant at $P < 0.05$.

Results

Linoleic acid affects Cx46 hemichannel currents

As previously described [47, 48], membrane depolarization of oocytes expressing rat Cx46 hemichannels induces slow outward currents that do not show inactivation (Fig. 1a).




Fig. 1 Cx46 hemichannels are inhibited by linoleic acid. **a** Representative whole-cell current records from *Xenopus* oocytes injected with Cx46 cRNA under control conditions. Oocytes were depolarized from -60 to $+60$ mV in steps of 10 mV for 15 s. At the end of each depolarizing step, the membrane was clamped to -60 mV for additional 10 s. **b** Same as **a** but after a 5 -min exposure to 100 μM linoleic acid (LA). **c** Average effect of 100 μM LA ($n=10$). **d** Average effect of 0.1 μM LA ($n=10$). Currents were normalized to that under

control conditions, at $+60$ mV, in the absence of LA. The symbols ** and *** correspond to $P<0.01$ and $P<0.001$ compared to the corresponding control values. **e** LA concentration dependence of Cx46 hemichannel currents. Results from ten oocytes were obtained at $+60$ mV and were normalized as described for **c**. The symbols *, **, and *** correspond to $P<0.05$, $P<0.01$, and $P<0.001$, respectively, compared to control condition, in the absence of LA. Average data in **c–e** are means \pm SEM

The maximum current at $+60$ mV was 5.5 ± 0.3 μA ($P<0.001$, $n=10$). Upon returning to the holding potential of -60 mV, small tail currents were observed (Fig. 1a). Addition of LA to the bath solution produced a biphasic response. At 0.1 μM LA, the currents increased $\sim 15\%$ (Fig. 1d), whereas at concentrations equal to or greater than 100 μM LA, both outward and tail currents were significantly reduced (Fig. 1b). With 100 μM LA, the current was reduced by $\sim 40\%$ (Fig. 1e), with $>90\%$ of the inhibition within 2 min of exposure (not shown). The inhibitory effect of LA was sustained for at least 30 min (Fig. 2a). To study the reversibility of the effect, we exposed oocytes for 5 min to

100 μM LA and then washed them for 5 min with ND96 without LA. The current, reduced 45% by LA, recovered to $\sim 90\%$ of the control value after washing (Fig. 2b). In four of the 12 experiments, the hemichannel current after wash actually increased to values similar to those elicited by 0.1 μM LA (not shown). The simplest explanation for this phenomenon and the dose–response effect in Fig. 1e is that LA has at least two mechanisms of action, one with higher affinity (that increases hemichannel currents) and other with lower affinity (that decreases hemichannel currents). Fits to the Boltzmann equation of the current data before and after 0.1 or 100 μM LA did not show changes in the voltage




Fig. 2 The linoleic acid effect is sustained and reversible. **a** Rapid and stable current block by 100 μ M LA. Oocytes were exposed to 100 μ M LA for 5, 15, or 30 min, and hemichannel currents were measured. Data were normalized and expressed as described in the legend to Fig. 1c. The solid lines are the fits of the Boltzmann equation to the data ($n=3$ each). **b** Reversibility of the inhibition by LA. Oocytes

expressing Cx46 hemichannels ($n=12$) were exposed to LA for 5 min (100 μ M LA), and hemichannel currents were recorded. Currents were measured again after washing without LA for 5 min (*Wash*). Currents were normalized to control values. A $P<0.001$ compared to control is denoted by **

dependency of Cx46 hemichannels, suggesting that LA modifies the open probability and/or the hemichannel conductance. In summary, LA has a biphasic effect, with a current increase at lower concentration and a decrease at higher concentrations. The inhibitory effect is complete in approximately 2 min and is reversible.

The inhibitory LA effect does not depend on an increase in intracellular free- $[Ca^{2+}]$ or PKC-mediated phosphorylation

Since LA can increase intracellular $[Ca^{2+}]$ [17] and also activate protein kinase C [31], we tested the effects of the Ca^{2+} chelator BAPTA-AM (100 μ M) and the PKC inhibitor

calphostin C (1 μ M) on the LA effect. Oocytes were pre-incubated for 1 h with one of these agents, briefly washed, exposed to 100 μ M LA for 5 min, and then Cx46 hemichannel currents were recorded. Neither BAPTA-AM nor calphostin C by themselves affected Cx46 hemichannel currents (data not shown, $n=6$), nor reduced the inhibitory effect of 100 μ M LA (Fig. 3). If anything, the LA effect was increased at the more positive clamping voltages. We did not examine further the increase in the response to LA, but one possibility is that BAPTA and calphostin C block the stimulatory effect elicited by lower LA concentrations (see Fig. 1, panels d and e). Nevertheless, the data strongly suggest that intracellular $[Ca^{2+}]$ or PKC-




Fig. 3 The effect of linoleic acid is not due to PKC- or Ca^{2+} -activated pathways. **a** Response to LA in oocytes incubated with 100 μ M BAPTA-AM for 1 h. Immediately after removing the Ca^{2+} chelator, the response to a 5-min exposure to 100 μ M LA was recorded and compared to control data from oocytes not treated with BAPTA-AM.

b Response to 100 μ M LA in oocytes incubated with 1 μ M calphostin C (*Calphos*) for 1 h. Immediately after removing the PKC blocker, the response to a 5-min exposure to 100 μ M LA was recorded and compared to control data from oocytes not treated with calphostin C. See legend to Fig. 1 for additional details

mediated phosphorylation is not necessary for the LA inhibitory effect.

Unsaturated fatty acids other than linoleic acid also inhibit Cx46 hemichannels

Oleamide and certain derivatives need specific chemical features, i.e., number and positions of double bonds, to affect GJC expressed in rat astrocytes [7]. Therefore, we tested the effects of fatty acids with different number of double bonds on Cx46 hemichannel currents. We used 100 μ M of each fatty acid and compared their effects to that of LA (18 carbons, two double bonds) at the same concentration. The choice of 100 μ M is based on concentrations of free C18:2 fatty acids of 50–100 μ M reported in human plasma [20, 28, 30]. The best Cx46 hemichannel inhibitor was arachidonic acid (AA) which is a polyunsaturated fatty acid with four double bonds and 20 carbons. The current reduction with AA was >70% (Fig. 4a). Oleic acid (OA), with one double bond and 18 carbons, reduced the Cx46 current by 20% (Fig. 4a). Palmitic acid (PA), with 16 carbons and no double bonds, had no effect on the current (Fig. 4a). These results suggest that the inhibition by fatty acids depends on the presence of double bonds and seems to be proportional to the degree of unsaturation.

A common characteristic between OA, LA, and AA is that all of them have a double bond at position 9, which seems to be important in the GJC inhibition by oleamide [7]. To test the importance of the double bond at position 9, we determined the effects of 100 μ M of 9-thiastearic acid (9-Th, C₁₇H₃₄O₂S) and 9-nitrooleate (9-NO, C₁₈H₃₃NO₄). The former has a sulfur atom at position 9, and the latter has a NO₂ at position 9, with the double bond not fully hydrogenated. The Cx46 hemichannel current was not affected by 5-min incubation with either of these fatty

acids (Fig. 4b). These results suggest that an unblocked double bond at carbon 9 is essential for the effect of fatty acids on Cx46 hemichannels.

Impermeable LA has no effects on hemichannel currents

Fatty acids are present in the inner and outer leaflets of the membrane bilayer because of fast flip-flop of the protonated species [29]. To test whether LA incorporation into the membrane inner leaflet is necessary for the fatty acid inhibitory effect, we studied the hemichannel response to LA attached to coenzyme A (LA-CoA, MW 1,029). LA-CoA is inserted in the outer leaflet of the plasma membrane, but as other CoA-linked fatty acids, it is not expected to flip to the internal monolayer [9]. The addition of 100 μ M LA-CoA to the bath solution for 5 min had no effect on Cx46 hemichannel currents (Fig. 5a). However, when LA-CoA was microinjected into the oocytes to a final estimated concentration of ca. 250 μ M, it did not decrease, but increased Cx46 hemichannel currents (Fig. 5b). In contrast, a decrease was observed when LA was microinjected to reach ca. 250 μ M (Fig. 5b).

LA does not affect Cx46 GJC currents

The data presented earlier showed that LA at 0.1 μ M activates and at 100 μ M inhibits Cx46 hemichannel currents. Here, we tested the effects of both concentrations on GJC currents in paired oocytes expressing Cx46. Under control conditions, Cx46 GJC exhibit currents of about 300 nA (283 \pm 63 nA at V_j -100 mV, and 361 \pm 36 nA at V_j \pm 100 mV), with a clear relaxation at V_j over 80 mV (Fig. 6a). A 5-min exposure to 0.1 or 100 μ M LA did not affect the GJC currents (data at +60 mV are shown in Fig. 6b). These results indicate that LA, at the same concentration and exposure time that affects




Fig. 4 Inhibition of Cx46 hemichannel currents by other unsaturated fatty acids. **a** Dependence of the inhibitory effect on the number of unsaturated bonds. The effects of exposure to palmitic acid (PA, $n=5$), oleic acid (OA, $n=5$), linoleic acid (LA, $n=6$), or arachidonic acid (AA, $n=6$) for 5 min at 100 μ M were determined. Protocol as described for Fig. 1d. Data are presented as means \pm SEM. The labeling of the bars shows, for each fatty acid, the number of carbons followed by the

number of double bonds; e.g., AA 20:4 for arachidonic acid, with 20 carbons and four double bonds. **b** Importance of the double bond at position 9. The experimental protocol was identical to that in **a**, but the fatty acids used were 9-thiastearic acid (9-Th) and 9-nitrooleate (9-NO). No statistically significant effects on the Cx46 hemichannel current were detected ($n=7$ for each compound). **c** Chemical structures of 9-thiastearic acid and 9-nitrooleate




Fig. 5 Microinjection of linoleic acid-CoA increases Cx46 hemichannel currents. **a** Current-voltage relationship of Cx46 hemichannel currents under control conditions and after a 5-min exposure to 100 μ M linoleic acid attached to coenzyme A (*LA-CoA*) or LA alone added to the extracellular solution ($n=8$ for each group). **b** Current-voltage

relationship of hemichannel currents under control conditions and 5 min after microinjection of LA-CoA ($n=15$) or LA ($n=14$) to reach final concentrations of ca. 250 μ M. Intracellular concentrations were estimated assuming a 1- μ l oocyte volume. Data are presented as means \pm SEM of currents normalized to the control value at +60 mV

hemichannel currents, does not alter the electrical properties of GJC.

Discussion

Mice expressing certain Cx46 mutants [44] and a Cx46 knockout [22] develop cataracts, clearly pointing to an essential role of Cx46 in maintaining lens transparency. Cataracts are frequent in diabetes [8, 56], and their incidence increases with age [34]; however, few studies have correlated cataract formation with potential alterations in Cx46 function induced by aging or diabetes. The studies available focused mostly on Cx46 modifications due to increases in free radicals [4, 47] and changes in phosphorylation state [19], but there are no studies on the effects of fatty acids on lens connexins, even though high concentrations of unsaturated fatty acids are toxic to human lens epithelial cells [27, 37]. In the present studies, we show the effect of physiological levels of unsaturated fatty acids on Cx46 hemichannels and GJCs. We focused our work on LA because it is present at the plasma membrane of lens epithelial cells and elevations of its concentration in the lens produce or accelerate cataract formation [37]. LA had a biphasic effect, increasing Cx46-hemichannel currents at low concentrations and decreasing the currents at higher concentrations. This is consistent with a high-affinity stimulatory effect and a low-affinity inhibitory effect. Dual effects of arachidonic acid on plasma-membrane Ca^{2+} -ATPase [42] and of oleic acid on gap-junction coupling in A7r5 cells [25] have also been reported.

In order to investigate the mechanism of action of LA, we evaluated the roles of increases in intracellular free- $[\text{Ca}^{2+}]$ and stimulation of PKC, two known responses to LA in

other systems [17, 31]. We found no involvement of either signaling pathway in the hemichannel inhibitory response to LA. Modulation of BKCa^{2+} channels [13], voltage-gated K^{+} channels [18], TRP channels [39], and voltage-gated




Fig. 6 Linoleic acid does not affect GJC currents. **a** Representative transjunctional current records from paired oocytes under control conditions. Both cells of the pair were clamped at -40 mV, and junctional currents were measured after changing the cell-membrane voltage of one cell to values between -140 and $+60$ mV, in 20-mV steps. **b** Lack of effect of LA on transjunctional conductance. GJC currents measured at +60 mV in the presence of 0.1 or 100 μ M LA were normalized to the current at +60 mV under control conditions. No differences were observed when 0.1 μ M or 100 μ M LA were used. Data are means \pm SEM ($n=6$ for each concentration used)

Ca^{2+} channels [49] by fatty acids has been reported, and in most cases, it has been suggested that there is a direct interaction between lipid and ion channel [14, 15, 49]. For example, arachidonic acid appears to bind to Thr250 and Val275, in the pore of Ca^{2+} -activated K^{+} channels [24].

Polyunsaturated fatty acids increase BK channel activity by shifting the voltage dependence of the open probability [15]. In this study, the magnitudes of the effects of OA, LA, and AA were in the same order as our inhibitory results on Cx46 hemichannels. *Cis*-unsaturated fatty acids disorder the bilayer interior and order the head-group region [31], and therefore, the effects on membrane proteins could be mediated by changes in the biophysical properties of the phospholipid membrane. Several facts argue against this interpretation of our results. In a study using 100 μM concentrations, the oil/water partition coefficients were $\text{PA} > \text{OA} > \text{LA} > \text{AA}$ [1], i.e., the opposite sequence to their inhibition of hemichannel conductance. In addition, in the BK channel study quoted above [15], there was no correlation between membrane fluidity and BK channel activation in response to the fatty acids. In sum, our results do not support an effect of LA and other unsaturated fatty acids mediated by changes in the properties of the plasma membrane. Nevertheless, additional experiments will be needed to determine whether the effects of fatty acids on Cx46 are due to direct binding to the hemichannels or are mediated by more complex signaling mechanisms, as suggested by the LA-CoA results discussed below. In addition, since Cx46 and Cx50 are the main lens connexin [35] and they form heteromeric hemichannels [26], further studies will be needed to determine whether hemichannels containing Cx50 are also affected by unsaturated fatty acids.

The absence of response to extracellular LA-CoA, not expected to flip across to the internal monolayer and therefore not to cross lipid bilayers [9], suggests that fatty acids need to access the inner leaflet of the plasma membrane and/or enter the intracellular compartment to affect hemichannels. Since the intracellular injection of sufficient LA-CoA to reach inhibitory concentrations produces an increase in Cx46 hemichannel currents, it is obvious that the mechanisms of stimulation and inhibition by LA are distinct. The current-activation site is accessible from the intracellular compartment to both LA and LA-CoA, whereas the inhibitory site is only accessible to LA, suggesting that it is located in a region of the hemichannel pore inaccessible to LA-CoA or that the signaling mechanism involved requires crossing of LA through an organelle membrane. Understanding of the complex mechanisms of the fatty acid effects on hemichannels will require additional studies.

Hemichannels are affected by changes in membrane phospholipid composition [36]. GJs formed by Cx26 or Cx32 associate preferentially with certain phospholipids;

phosphatidylcholine does not associate with Cx26 hemichannels, whereas phosphatidylserine does not associate with Cx32 hemichannels [11, 36]. In contrast, phosphatidylcholine and phosphatidyl serine are associated with both isoforms in junctional plaques [36]. The concept of specific lipid–connexin interaction is also supported by Boger and co-workers [7], who found that molecules derived from the fatty acid primary amide oleamide affect GJs when they contain 16–24 carbon atoms, a hydrophobic methyl terminus, a polarized carboxy-terminal and a *cis* double bond at position 9. In agreement with these observations, our results with fatty acids containing varied number of double bonds, with and without an unsaturated *cis* double bond at position 9, support the idea that the inhibitory effect of the fatty acids requires the unsaturated *cis* double bond at position 9 and is increased by the degree of unsaturation. As mentioned above, some phospholipids interact differentially with hemichannel or GJs formed by Cx26 and Cx32 [36]. Our results showing that LA affects hemichannels, but not the GJs formed by Cx46, extend previous studies to a lens connexin and show differences in functional responses between GJs and hemichannels. We have not determined whether the selective effect on hemichannels vs. GJs is due to differences in affinity for LA of hemichannels and GJs. However, since LA is present on both leaflets of the plasma membrane, the absence of effect on GJs may be due to structural differences between hemichannels and GJs.

Pathophysiological significance

Unsaturated fatty acids modulate a wide range of cellular processes [52], and in the lens they induce or accelerate cataract formation by unknown mechanisms [21, 27, 33, 37, 40]. The cytotoxic effects of fatty acids on lens fibers depend on unsaturated fatty acid uptake by the cells and increase with the fatty acid-to-albumin ratio [27, 57]. In aqueous humor, the concentration of albumin is very low compared to that in plasma [27], and increases in the fatty acid-to-albumin ratio occur in a number of conditions associated with cataracts, including diabetes, lipid disorders, nephritic syndrome, and liver diseases [see 27]. Under these conditions, unsaturated fatty acid damage of the lens fibers can occur, particularly in the elderly, where there is a significant loss of the blood-aqueous humor barrier [see 27].

The block of Cx46 hemichannel currents by LA and other unsaturated fatty acids at concentrations found in the lens, without affecting Cx46 GJs, may contribute to the induction of cataracts by: (a) reducing the uptake of physiological molecules from the extracellular space, as has been proposed for astrocytes [46], (b) reducing the release of signaling molecules, as it has been shown in many cell types [50], and/or (c) diminishing cell-to-cell

communication. It has been proposed that the activity of Cx46 hemichannels mediates GJC formation [3], and therefore, reduced hemichannel activity may decrease the number of GJCs.

Acknowledgments This work was supported by the National Institutes of Health grants GM068586, GM79629, and R01GM079629-03S1; Texas Advanced Research Program grant 010674-0046-2007; Fondecyt de Iniciación 11080061; and Proyecto Interno Universidad del Desarrollo 23.400.012.

References

1. Anel A, Richieri GV, Kleinfeld AM (1993) Membrane partition of fatty acids and inhibition of T cell function. *Biochemistry* 32:530–536
2. Bao X, Altenberg GA, Reuss L (2004) Mechanism of regulation of the gap junction protein connexin 43 by protein kinase C-mediated phosphorylation. *Am J Physiol Cell Physiol* 286:C647–C654
3. Beahm DL, Hall JE (2004) Opening hemichannels in nonjunctional membrane stimulates gap junction formation. *Biophys J* 86:781–796
4. Berthoud VM, Beyer EC (2009) Oxidative stress, lens gap junctions, and cataracts. *Antioxid Redox Signal* 11:339–353
5. Besancon E, Guo S, Lok J, Tymianski M, Lo EH (2008) Beyond NMDA and AMPA glutamate receptors: emerging mechanisms for ionic imbalance and cell death in stroke. *Trends Pharmacol Sci* 29:268–275
6. Beyer EC, Kistler J, Paul DL, Goodenough DA (1989) Antisera directed against connexin43 peptides react with a 43-kD protein localized to gap junctions in myocardium and other tissues. *J Cell Biol* 108:595–605
7. Boger DL, Patterson JE, Guan X, Cravatt BF, Lerner RA, Gilula NB (1998) Chemical requirements for inhibition of gap junction communication by the biologically active lipid oleamide. *Proc Natl Acad Sci USA* 95:4810–4815
8. Borenshtein D, Ofri R, Werman M, Stark A, Tritschler HJ, Moeller W, Madar Z (2001) Cataract development in diabetic sand rats treated with alpha-lipoic acid and its gamma-linolenic acid conjugate. *Diabetes/Metab Res Rev* 17:44–50
9. Boylan JG, Hamilton JA (1992) Interactions of acyl-coenzyme A with phosphatidylcholine bilayers and serum albumin. *Biochemistry* 31:557–567
10. Bruzzone S, Guida L, Zocchi E, Franco L, De Flora A (2001) Connexin 43 hemi channels mediate Ca^{2+} -regulated transmembrane NAD^+ fluxes in intact cells. *FASEB J* 15:10–12
11. Cascio M (2005) Connexins and their environment: effects of lipids composition on ion channels. *Biochim Biophys Acta* 1711:142–153
12. Cherian PP, Siller-Jackson AJ, Gu S, Wang X, Bonewald LF, Sprague E, Jiang JX (2005) Mechanical strain opens connexin 43 hemichannels in osteocytes: a novel mechanism for the release of prostaglandin. *Mol Biol Cell* 16:3100–3106
13. Clarke AL, Petrou S, Walsh JV Jr, Singer JJ (2002) Modulation of BK(Ca) channel activity by fatty acids: structural requirements and mechanism of action. *Am J Physiol Cell Physiol* 283:C1441–C1453
14. Clarke AL, Petrou S, Walsh JV Jr, Singer JJ (2003) Site of action of fatty acids and other charged lipids on BKCa channels from arterial smooth muscle cells. *Am J Physiol Cell Physiol* 284:C607–C619
15. Denson DD, Wang X, Worrell RT, Eaton DC (2000) Effects of fatty acids on BK channels in GH(3) cells. *Am J Physiol Cell Physiol* 279:C1211–C1219
16. Ebihara L, Tong JJ, Vertel B, White TW, Chen TL (2011) Properties of connexin 46 hemichannels in dissociated lens fiber cells. *Invest Ophthalmol Vis Sci* 52:882–889
17. El-Yassimi A, Hichami A, Besnard P, Khan NA (2008) Linoleic acid induces calcium signaling, Src kinase phosphorylation, and neurotransmitter release in mouse CD36-positive gustatory cells. *J Biol Chem* 283:12949–12959
18. Feng DD, Luo Z, Roh SG, Hernandez M, Tawadros N, Keating DJ, Chen C (2006) Reduction in voltage-gated K^+ currents in primary cultured rat pancreatic beta-cells by linoleic acids. *Endocrinology* 147:674–682
19. Fleschner CR (2006) Connexin 46 and connexin 50 in selenite cataract. *Ophthalmic Res* 38:24–28
20. Fraser DA, Thoen J, Rustan AC, Førre O, Kjeldsen-Kragh J (1999) Changes in plasma free fatty acid concentrations in rheumatoid arthritis patients during fasting and their effects upon T-lymphocyte proliferation. *Rheumatol Oxf* 38:948–952
21. Glanz D, Lennarz B, Glaesser D (2006) Fatty acid cytotoxicity to organ-cultured bovine lenses. *Ophthalmic Res* 38:62–65
22. Gong X, Li E, Klier G, Huang Q, Wu Y, Lei H, Kumar NM, Horwitz J, Gilula NB (1997) Disruption of alpha3 connexin gene leads to proteolysis and cataractogenesis in mice. *Cell* 91:833–843
23. Goodenough DA (1979) Lens gap junctions: a structural hypothesis for nonregulated low-resistance intercellular pathways. *Invest Ophthalmol Vis Sci* 18:1104–1122
24. Hamilton KL, Syme CA, Devor DC (2003) Molecular localization of the inhibitory arachidonic acid binding site to the pore of hIK1. *J Biol Chem* 278:16690–16697
25. Hirschi KK, Minnich BN, Moore LK, Burt JM (1993) Oleic acid differentially affects gap junction-mediated communication in heart and vascular smooth muscle cells. *Am J Physiol Cell Physiol* 265:C1517–C1526
26. Hopperstad MG, Srinivas M, Spray DC (2000) Properties of gap junction channels formed by Cx46 alone and in combination with Cx50. *Biophys J* 79:1954–1966
27. Iwig M, Glaesser D, Fass U, Struck HG (2004) Fatty acid cytotoxicity to human lens epithelial cells. *Exp Eye Res* 79:689–704
28. Jahn CE, Leiss O, von Bergmann K (1983) Lipid composition of human aqueous humor. *Ophthalmic Res* 15:220–224
29. Kamp F, Westerhoff HV, Hamilton JA (1992) pH gradients across phospholipid membranes caused by fast flip-flop of un-ionized fatty acids. *Proc Natl Acad Sci USA* 89:11367–11370
30. Kangani CO, Kelley DE, DeLany JP (2008) New method for GC/FID and GC-C-IRMS analysis of plasma free fatty acid concentration and isotopic enrichment. *J Chromatogr B* 873:95–101
31. Kim MH, Kim MO, Kim YH, Kim JS, Han HJ (2009) Linoleic acid induces mouse embryonic stem cell proliferation via Ca^{2+} /PKC, PI3K/Akt, and MAPKs. *Cell Physiol Biochem* 23:53–64
32. Kistler J, Kirkland B, Bullivant S (1985) Identification of a 70,000-D protein in lens membrane junctional domains. *J Cell Biol* 101:28–35
33. Kojima K, Okochi Y, Yagi K (1968) The opacity of the rat lens caused by fatty acid peroxide. *Nippon Ganka Gakkai Zasshi* 72:1733–1739
34. Kronfeld PC (1953) Eye changes due to advanced age. *Ill Med J* 103:104–107
35. Lo WK, Shaw AP, Takemoto LJ, Grossniklaus HE, Tigges M (1996) Gap junction structures and distribution patterns of immunoreactive connexins 46 and 50 in lens regrowths of Rhesus monkeys. *Exp Eye Res* 62:171–180
36. Locke D, Harris AL (2009) Connexin channels and phospholipids: association and modulation. *BMC Biol* 17:7–52

37. Lu M, Taylor A, Chylack LT Jr, Rogers G, Hankinson SE, Willett WC, Jacques PF (2005) Dietary fat intake and early age-related lens opacities. *Am J Clin Nutr* 81:773–779
38. Mathias RT, Rae JL, Baldo GJ (1997) Physiological properties of the normal lens. *Physiol Rev* 77:21–50
39. Matta JA, Miyaresm RL, Ahern GP (2007) TRPV1 is a novel target for omega-3 polyunsaturated fatty acids. *J Physiol* 578:397–411
40. Nguyen N, Glanz D, Glaesser D (2000) Fatty acid cytotoxicity to bovine lens epithelial cells: investigations on cell viability, ecto-ATPase, Na(+), K(+)-ATPase and intracellular sodium concentrations. *Exp Eye Res* 71:405–413
41. Nojima H, Ohba Y, Kita Y (2007) Oleamide derivatives are prototypical anti-metastasis drugs that act by inhibiting Connexin 26. *Curr Drug Saf* 2:204–211
42. Oliveira VH, Nascimento KS, Freire MM, Moreira OC, Scofano HM, Barrabin H, Mignaco JA (2008) Mechanism of modulation of the plasma membrane Ca(2+)-ATPase by arachidonic acid. *Prostaglandins Other Lipid Mediat* 87:47–53
43. Pal JD, Berthoud VM, Beyer EC, Mackay D, Shiels A, Ebihara L (1999) Molecular mechanism underlying a Cx50-linked congenital cataract. *Am J Physiol Cell Physiol* 276:C1443–C1446
44. Pal JD, Liu X, Mackay D, Shiels A, Berthoud VM, Beyer EC, Ebihara L (2000) Connexin46 mutations linked to congenital cataract show loss of gap junction channel function. *Am J Physiol Cell Physiol* 279:C596–C602
45. Paul DL, Ebihara L, Takemoto LJ, Swenson KI, Goodenough DA (1991) Connexin46, a novel lens gap junction protein, induces voltage-gated currents in nonjunctional plasma membrane of *Xenopus* oocytes. *J Cell Biol* 115:1077–1089
46. Retamal MA, Froger N, Palacios-Prado N, Ezan P, Sáez PJ, Sáez JC, Giaume C (2007) Cx43 hemichannels and gap junction channels in astrocytes are regulated oppositely by proinflammatory cytokines released from activated microglia. *J Neurosci* 27:13781–13792
47. Retamal MA, Yin S, Altenberg GA, Reuss L (2009) Modulation of Cx46 hemichannels by nitric oxide. *Am J Physiol Cell Physiol* 296:C1356–C1363
48. Retamal MA, Yin S, Altenberg GA, Reuss L (2010) Voltage-dependent facilitation of Cx46 hemichannels. *Am J Physiol Cell Physiol* 298:C132–C139
49. Roberts-Crowley ML, Rittenhouse AR (2009) Arachidonic acid inhibition of L-type calcium (CaV1.3b) channels varies with accessory CaVbeta subunits. *J Gen Physiol* 133:387–403
50. Sáez JC, Schalper KA, Retamal MA, Orellana JA, Shoji KF, Bennett MV (2010) Cell membrane permeabilization *via* connexin hemichannels in living and dying cells. *Exp Cell Res* 316:2377–2389
51. Schalper KA, Palacios-Prado N, Retamal MA, Shoji KF, Martínez AD, Sáez JC (2008) Connexin hemichannel composition determines the FGF-1-induced membrane permeability and free $[Ca^{2+}]_i$ responses. *Mol Biol Cell* 19:3501–3513
52. Schmitz G, Ecker J (2008) The opposing effects of n-3 and n-6 fatty acids. *Prog Lipid Res* 47:147–155
53. Steinberg G, Slaton WH Jr, Howton DR, Mead JF (1956) Metabolism of essential fatty acids. IV. Incorporation of linoleate into arachidonic acid. *J Biol Chem* 220:257–264
54. Stout CE, Costantin JL, Naus CC, Charles AC (2002) Intercellular calcium signaling in astrocytes *via* ATP release through connexin hemichannels. *J Biol Chem* 277:10482–10488
55. Stridh MH, Tranberg M, Weber SG, Blomstrand F, Sandberg M (2008) Stimulated efflux of amino acids and glutathione from cultured hippocampal slices by omission of extracellular calcium: likely involvement of connexin hemichannels. *J Biol Chem* 283:10347–10356
56. Tan JS, Wang JJ, Mitchell P (2008) Influence of diabetes and cardiovascular disease on the long-term incidence of cataract: the Blue Mountains eye study. *Ophthalmic Epidemiol* 15:317–327
57. Trimborn M, Iwig M, Glanz D, Gruner M, Glaesser D (2000) Linoleic acid cytotoxicity to bovine lens epithelial cells: influence of albumin on linoleic acid uptake and cytotoxicity. *Ophthalmic Res* 32:87–93
58. Ye ZC, Wyeth MS, Baltan-Tekkok S, Ransom BR (2003) Functional hemichannels in astrocytes: a novel mechanism of glutamate release. *J Neurosci* 23:3588–3596