



Research Article

Association of Dual-Phase Computed Tomography Angiography and CT-Perfusion in Patients with Acute Ischemic Stroke Beyond the 6-Hour Window

Andrés Gallardo Vera^{1,3,6*}, Pablo M Lavados^{2,6}, Pablo Cox Vial^{3,4}, Camila de la Barra⁵, Gabriel Cavada⁶ and Verónica V Olavarria^{2,6}

¹Unidad de TeleACV, Hospital Digital, MINSAL, Chile

²Servicio de Neurología, Departamento de Neurología y Psiquiatría, Clínica Alemana de Santiago, Facultad de Medicina Clínica Alemana Universidad del Desarrollo, Santiago, Chile

³Unidad de Imagenología Compleja, Hospital Carlos Van Buren, Valparaíso, Chile

⁴Centro de Neurología Traslacional, Universidad de Valparaíso, Chile

⁵Departamento de Imagenología, Clínica Alemana de Santiago, Facultad de Medicina Clínica Alemana Universidad del Desarrollo, Santiago, Chile

⁶Unidad de Investigación y Ensayos Clínicos, Departamento Científico Docente, Clínica Alemana de Santiago, Santiago, Chile

ARTICLE INFO

Article history:

Received: 22 May, 2023

Accepted: 14 June, 2023

Published: 11 July, 2023

Keywords:

Dual-phase computed tomography angiography, CT-Perfusion, late window acute ischemic stroke

ABSTRACT

Background: There is controversy regarding the need to use advanced imaging to select candidates for thrombectomy in late window acute ischemic stroke (AIS). Hypoattenuation on CT angiography source images (CTA-SI) in arterial phase has been shown to be more sensitive than Alberta Stroke Program Early CT score (ASPECTS) of brain parenchyma to determine tissue at risk of ischemia. Our hypothesis is that the addition of a second acquisition at 35-50 seconds could complement the assessment of hypoperfused tissue that fails to receive flow through pial vessels. **Methods:** Patients with large vessel occlusion and 6-16 hours from symptom onset, admitted between August 2019 and July 2022 were evaluated with dual-phase CT angiography (CTA) and CT-perfusion. The association between automated CT-perfusion values and dual-phase CTA was assessed through a correlation coefficient. **Results:** Pearson's coefficient demonstrated a high correlation between ischemic core volume and delayed CTA-SI with an inverse association of -0.91 and between T_{max}≥6 sec volume and arterial CTA-SI with a value of -0.83. **Conclusion:** CTA-derived source images (CTA-SI) in two phases may be useful in the selection of patients with AIS presenting beyond the 6-hour window.

© 2023 Published by Progress in Neurobiology

1. Introduction

Acute Ischemic Stroke (AIS) is the leading cause of death and disability worldwide [1], and reperfusion therapies are very effective in reducing the number of patients who suffer a deterioration in their functional status due to AIS [2, 3]. There is relative consensus on the tendency to simplify imaging tests to select patients diagnosed with large vessel occlusion (LVO) in the early window and to dispense with multimodal neuroimaging [4]. However, the potential benefit of mechanical thrombectomy (MT) in patients with ≥6 hours of evolution depends on proper selection of slow progressors, which is directly related to a good leptomeningeal collaterality [5]. Perfusion studies have proven to be useful in predicting the volume of ischemic core and critical hypoperfusion [6], although they have the disadvantage of increasing

exposure to higher doses of contrast and radiation, in addition to requiring post-processing, which, even with automated software, prolongs the time required to decide on the therapeutic approach [7].

Perfusion studies are not widely available in primary stroke centers, which is a major limitation when it comes to being able to increase the time criteria for selection of patients susceptible to endovascular therapy that could be taken to comprehensive stroke centers. For this reason, selection methods based only on computed tomography (CT) in late-window patients are being explored [8]. The usual neuroimaging protocol for the study of AIS includes CT angiography (CTA) from the aortic arch to the vertex, which is usually performed in the arterial phase [9].

*Correspondence to: Andrés Gallardo Vera, M.D., Unidad de TeleACV, Hospital Digital, MINSAL, Huérfanos 670, Postal Code: 8320196, Unidad de Imagenología Compleja Hospital Carlos Van Buren, Valparaíso, Chile; ORCID: 0000-0002-7419-4771; E-mail: andres.gallardo@minsal.cl

Hypoattenuation in CT angiography source images (CTA-SI) in early arterial phase has been shown to be more sensitive than Alberta Stroke Program Early CT score (ASPECTS) of brain parenchyma to determine tissue at risk of ischemia, so it could be a good tool to determine viable salvageable tissue in an extended window [10, 11]. Our hypothesis is that the addition of a second acquisition at 35-50 seconds could complement the assessment of hypoperfused tissue that fails to receive flow through pial vessels.

2. Methodology

Our study is a retrospective analysis of a prospective cohort named Registro de Enfermedades Cerebrovasculares de Clínica Alemana de Santiago (RECCA), which registered consecutive adults patients under informed consent approved by the Institutional Ethics Committee. The initial registry included all adult patients with AIS admitted to Clínica Alemana de Santiago from August 2019 to July 2022. Study inclusion criteria were: i) patient with AIS between 6-16 hours from symptom onset, ii) NIHSS ≥ 6 , iii) LVO of anterior circulation, iv) admission CT-perfusion and CTA in arterial/delayed phase.

2.1. Exclusion Criteria

Incomplete data.

2.2 Image and Interpretation Protocol

The post-processing of CT-perfusion-DWI/ADC images was performed in an automated way through RAPID© software, which considers a candidate for extended window endovascular therapy those patients whose ischemic core determined by $CBF < 30\%$ was not greater than 50 ml, in $DWI \leq 70$ ml and the mismatch between $T_{max} > 6$ sec/ $CBF < 30\%$ volume ≥ 1.8 .

CTA from supra-aortic trunks to vertex was performed in arterial and delayed phase at 35-50 seconds. A vascular neurologist performed the estimation of CTA-SI ASPECTS in both phases at the time of data entry into the RECCA registry, which was done blind to the results of the perfusion study. A parallel evaluation of the CTA-SI was done by a neuroradiologist who was blind to both the clinical history and the results of the multimodal neuroimaging with automated processing. In case of a discrepancy, consensus could be reached by considering a third evaluation carried out by an external neuroradiologist. Mismatch was considered when the difference between arterial CTA-SI and delayed phase CTA-SI was ≥ 2 points (Figures 1 & 2).




FIGURE 1: Occlusion in M1 segment of the left middle cerebral artery. **A & B)** Correspond to the arterial phase. **C & D)** Correspond to the delayed phase with filling by leptomeningeal vessels in the M1-M2-M4 and M5 region of ASPECTS. Considering a significant mismatch, the patient underwent mechanical thrombectomy.




FIGURE 2: Occlusion in M1 segment of the left middle cerebral artery. **A & B)** In arterial phase with low CTA-SI ASPECTS, **C & D)** In delayed phase without significant changes.

2.3. Statistical Analysis

The normality of quantitative variables was evaluated through the Shapiro-Wilk test, using descriptive statistics to calculate the mean, standard deviation, median and interquartile range. Strength of association between continuous variables of the automated perfusion study and the CTA evaluation was assessed through Pearson's correlation coefficient. In addition, the association between categorical variables of the automated study and visual assessment was assessed using Fisher's exact test. To predict the outcome of a categorical variable of the perfusion study by logistic regression, all clinical and demographic characteristics at admission were used as independent variables, as well as the results of the CTA study. The data were processed with STATA version 16.0

3. Results

During the study period, 15 patients met the inclusion criteria. The mean age was 77.8 years (range 48-92 years), the median NIHSS score was 14 (IQR 10-19) and the median time from symptom onset to CT was 10.5 hours (Table 1). Arterial CTA-SI/delayed phase CTA-SI mismatch was present in 9 of the 15 patients, with no discrepancy between vascular neurologist and neuroradiologist. Pearson's coefficient demonstrated a high correlation between ischemic core volume and delayed CTA-SI with an inverse association of -0.91 $p < 0.0001$ (Figure 3), and between $T_{max} \geq 6$ sec volume and arterial CTA-SI with a value of -0.83 $p < 0.0001$ (Figure 4).

TABLE 1: Baseline clinical and imaging characteristics.

Variable	N	p50	mean	SD
Age	15	83	77.8	13.5
NIHSS admission	15	14	15	7.4
Time onset-CT	15	10.5	10.3	2.7
Core volume (CBF<30%)	15	27	56.5	68.5
Critical Hypoperfusion ($T_{max} > 6$ sec)	15	118	141.2	78.2
CTA-SI Arterial ASPECTS	15	6	6.1	1.3
CTA-SI Delayed ASPECTS	15	8	7.7	1.8

NIHSS: National Institutes of Health Stroke Scale; ASPECTS: Alberta Stroke Program Early CT Score; CBF: Cerebral Blood Flow; T_{max} : Time-to-maximum.




FIGURE 3: Scatter plot of Pearson`s correlation coefficient. A significant negative correlation between ischemic core volume (x-axis) and delayed CTA-SI (y-axis).




FIGURE 4: Scatter plot of Pearson`s correlation coefficient. A significant negative correlation between $T_{max} \geq 6sec$ (x-axis) volume and arterial CTA-SI (y-axis).

There was a significant association between patient selection under DEFUSE 3 (Endovascular Therapy Following Imaging Evaluation for Ischemic Stroke 3) criteria and the assessment of arterial CTA-SI/delayed CTA-SI mismatch with a p-value of 0.002. In addition, there was a direct relationship between the probability of being a candidate for MT by RAPID automated CT-perfusion criteria and a higher CTA-SI ASPECTS value in delayed phase, with a perfect association for CTA-SI ASPECTS values ≥ 8 .

4. Discussion

Hypoattenuation on CTA-SI has been evaluated qualitatively with CBF and CBV maps, demonstrating a higher correlation with CBF drop, suggesting a significant approximation to critically hypoperfused tissue [12]. Although, to date the association has only been made from CTA-SI in arterial time and there is no data when it is compared with the volumetric values offered by the automated software. Adequate assessment of the volume of hypoperfused and infarcted tissue is essential for the selection of slow progressors [13, 14], who are potential beneficiaries of late-window endovascular therapy [15]. However, according to our results, the assessment of dual-phase CTA seems to be a semiquantitative tool comparable to the multimodal study with validated metrics.

Delayed-phase CTA-SI showed a high approximation to what we consider nonviable tissue, with a 50th percentile on ASPECTS 8 appearing to be a possible upper limit for ischemic core. Moreover, the CTA-SI in the arterial phase presented a high correlation with what we determined by imaging consensus as critical hypoperfusion, which includes viable tissue. The mean difference between arterial and delayed CTA-SI was 2 points, so it could be hypothesized that a mismatch ≥ 2 would be a value to validate in future studies of larger sample size. Therefore, these results provide information in the direction of the feasibility of choosing patients in extended window with imaging protocols with greater availability, shorter acquisition time and lower irradiation. In addition, it is possible that dispensing with CT-perfusion could increase the number of patients who can benefit from TM, since there is growing evidence that advanced imaging studies lead to overselection [16].

Although a clinical trial recently demonstrated that identifying some degree of collateral flow in the arterial phase makes it possible to identify intermediate progressors that benefit from mechanical thrombectomy [17], it is still essential to identify slow progressors when a transfer of more than 1 hour to the stroke center is required. Our study has some limitations. First, because of its retrospective nature, we cannot exclude a certain selection bias. Second, the small sample size decreases the power of the study and increases the margin of error.

In conclusion, dual-phase CTA could be used similar to CT-perfusion and derive information on salvageable and non-salvageable tissue. It could also be used to select patients eligible for thrombectomy in the late time window, however, because of study limitations, these findings should be interpreted as preliminary and require replication.

REFERENCES

- [1] GBD 2017 DALYs, HALE Collaborators "Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017." *Lancet*, vol. 392, no. 10159, pp. 1859-1922, 2018. View at: [Publisher Site](#) | [PubMed](#)
- [2] Mayank Goyal, Bijoy K Menon, Wim H van Zwam, et al. "Endovascular thrombectomy after large-vessel ischaemic stroke: a meta-analysis of individual patient data from five randomised trials." *Lancet*, vol. 387, no. 10029, pp. 1723-1731. View at: [Publisher Site](#) | [PubMed](#)
- [3] Wondwossen G Tekle, Saqib A Chaudhry, Zara Fatima, et al. "Intravenous Thrombolysis in Expanded Time Window (3-4.5 hours) in General Practice with Concurrent Availability of Endovascular Treatment." *J Vasc Interv Neurol*, vol. 5, no. 1, pp. 22-26, 2012. View at: [PubMed](#)
- [4] Manuel Requena, Natalia Pérez de la Ossa, Sonia Abilleira, et al. "Predictors of endovascular treatment among stroke codes activated within 6 hours from symptom onset." *Stroke*, vol. 49, no. 9, pp. 2116-2121, 2018. View at: [Publisher Site](#) | [PubMed](#)
- [5] Igor Pagiola, Olivier Chassin, Sophie Gallas, et al. "Extending the stroke treatment window beyond DAWN in patients with very slow progressor type collaterals: How far can we go?" *J Cerebrovasc Endovasc Neurosurg*, vol. 23, no. 4, pp. 354-358, 2021. View at: [Publisher Site](#) | [PubMed](#)
- [6] Gregory W Albers, Michael P Marks, Stephanie Kemp, et al. "Thrombectomy for stroke at 6 to 16 hours with selection by perfusion imaging." *N Engl J Med*, vol. 378, no. 8, pp. 708-718, 2018. View at: [Publisher Site](#) | [PubMed](#)
- [7] Jeremy J Heit, Max Wintermark "Perfusion Computed Tomography for the Evaluation of Acute Ischemic Stroke: Strengths and Pitfalls." *Stroke*, vol. 47, no. 4, pp. 1153-1158, 2016. View at: [Publisher Site](#) | [PubMed](#)
- [8] Thanh N Nguyen, Mohamad Abdalkader, Simon Nagel, et al. "Noncontrast computed tomography vs computed tomography perfusion or magnetic resonance imaging selection in late presentation of stroke with large-vessel occlusion." *JAMA Neurol*, vol. 79, no. 1, pp. 22-31, 2022. View at: [Publisher Site](#) | [PubMed](#)
- [9] Bijoy K Menon, Mayank Goyal "Imaging Paradigms in Acute Ischemic Stroke: A Pragmatic Evidence-based Approach." *Radiology*, vol. 277, no. 1, pp. 7-12. View at: [Publisher Site](#) | [PubMed](#)
- [10] Jung-Soo Park, Jong Myong Lee, Hyo-Sung Kwak, et al. "Predictive value of CT angiography source image ASPECTS in patients with anterior circulation acute ischemic stroke after endovascular treatment: ultimate infarct size and clinical outcome." *J Neurointerv Surg*, vol. 11, no. 4, pp. 342-346. View at: [Publisher Site](#) | [PubMed](#)
- [11] Shelagh B Coutts, Michael H Lev, Michael Eliasziw, et al. "ASPECTS on CTA source images versus unenhanced CT: added value in predicting final infarct extent and clinical outcome." *Stroke*, vol. 35, no. 11, pp. 2472-2476, 2004. View at: [Publisher Site](#) | [PubMed](#)
- [12] M Sharma, A J Fox, S Symons et al. "CT angiographic source images: flow- or volume-weighted?" *AJNR Am J Neuroradiol*, vol. 32, no. 2, pp. 359-364, 2011. View at: [Publisher Site](#) | [PubMed](#)

- [13] Marcelo Rocha, Shashvat Desai, Jiyeon Son, et al. "Clinical characteristics of fast and slow progressors of infarct growth in anterior circulation large vessel occlusion stroke." *J Cereb Blood Flow Metab*, vol. 41, no. 7, pp. 271678X211015068, 2021. View at: [Publisher Site](#) | [PubMed](#)
- [14] Marcelo Rocha, Tudor G Jovin "Fast Versus Slow Progressors of Infarct Growth in Large Vessel Occlusion Stroke: Clinical and Research Implications." *Stroke*, vol. 48, no. 9, pp. 2621-2627, 2017. View at: [Publisher Site](#) | [PubMed](#)
- [15] Raul G Nogueira, Ashutosh P Jadhav, Diogo C Haussen, et al. "Thrombectomy 6 to 24 hours after stroke with a mismatch between deficit and infarct." *N Engl J Med*, vol. 378, no. 1, pp. 11-21, 2018. View at: [Publisher Site](#) | [PubMed](#)
- [16] Guilherme B F Porto, Ching-Jen Chen, Sami Al Kasab, et al. "Association of Noncontrast Computed Tomography and Perfusion Modalities With Outcomes in Patients Undergoing Late-Window Stroke Thrombectomy." *JAMA Netw Open*, vol. 5, no. 11, pp. e2241291, 2022. View at: [Publisher Site](#) | [PubMed](#)
- [17] Susanne G H Olthuis, F Anne V Pirson, Florentina M E Pinckaers, et al. "Endovascular treatment versus no endovascular treatment after 6-24 h in patients with ischaemic stroke and collateral flow on CT angiography (MR CLEAN-LATE) in the Netherlands: a multicentre, open-label, blinded-endpoint, randomised, controlled, phase 3 trial." *Lancet*, vol. 401, no. 10385, pp. 1371-1380, 2023. View at: [Publisher Site](#) | [PubMed](#)